DOE/ER/41014-10-N97

# Non-$1/m_b^n$ Power Suppressed Contributions to Inclusive $B \to X_s\ l^+l^-$ Decays

Jiunn-Wei Chen, Gautam Rupak and Martin J. Savage*
*Department of Physics, University of Washington,*
*Seattle, WA 98195-1560.*

arXiv:hep-ph/9705219v1 2 May 1997

## Abstract

We compute non-perturbative contributions to $B \to X_s\ l^+l^-$ that are not explicitly suppressed by powers of the $b$-quark mass. They are proportional to $\langle B|\overline{h}\sigma\cdot Gh|B\rangle$ and arise from an interference between the free-quark amplitude and higher order terms in the matrix element of the four-quark operator $\overline{s}^\alpha\gamma^\mu(1-\gamma_5)c_\alpha\ \overline{c}^\beta\gamma_\mu(1-\gamma_5)b_\beta$. This correction is found to be small over most of the dalitz plot except near the charm threshold. Unfortunately, the perturbative computation we have performed is invalid near charm threshold and we do not except to see the structure found at lowest order reproduced in the data. We conclude that these non-perturbative contributions do not significantly modify the previous analysis of $B \to X_s\ l^+l^-$.

April 1997

*savage@phys.washington.edu

Rare decays of B mesons such as $B \to X_s\gamma$ and $B \to X_s\ l^+l^-$ provide a probe of physics at distance scales much smaller than the compton wavelength of the b-quark. The rates for these processes vanish at tree-level in the standard model of electroweak interactions but proceed at one-loop and beyond. Their GIM "suppressed" amplitudes are dominated by the loop contributions from the top quark. As such, $B \to X_s\gamma$ and $B \to X_s\ l^+l^-$ are ideal candidates for observing physics beyond that of the standard model, e.g. an extended Higgs sector. However, in order to use them to identify or even claim a hint of new physics one must have a robust prediction for decay rates and/or other observables in the standard model alone. Both the inclusive $B \to X_s\gamma$ (with cuts on the photon energy $E_\gamma$) and some exclusive radiative decay modes such as $B \to K^*\gamma$ have been observed. As yet neither inclusive nor exclusive decays $B \to X_s\ l^+l^-$ have been observed, however, the CLEO [1] upper limit of $1.6 \times 10^{-5}$ on the $\overline{B}^0 \to \overline{K}^{*0}e^+e^-$ branching ratio is within a factor of $\sim 3$ of a standard model estimate. As theoretical predictions for exclusive decay widths are notoriously difficult to obtain it is the inclusive decays that one is presently looking toward to make a meaningful comparison between theory and experiment.

The heavy quark limit of QCD leads to great simplification in the theoretical analysis of inclusive decays of B mesons [2]. In particular, the inclusive decays of B mesons can be described using an operator-product expansion since the characteristic momentum of the process $\sim m_b$ is large compared to the scale of strong interactions $\sim \Lambda_{\rm QCD}$. A systematic expansion of decay observables, such as the rate and differential spectra can be performed in powers of $\Lambda_{\rm QCD}/m_b$ and the strong coupling constant $\alpha_s(\mu)$. There are three non-perturbative matrix elements, $\overline{\Lambda}$, $\lambda_1$ and $\lambda_2$, [3] that appear in expressions for such observables up to order $1/m_b^2$ in the expansion.

The analysis of the inclusive decays $B \to X_s\gamma$ [4,5] and $B \to X_s\ l^+l^-$ [4] to $\mathcal{O}\left(1/m_b^2\right)$ is slightly different to the analogous computations performed for inclusive semileptonic decays. The difference lies in the fact that the semileptonic decays proceed at tree-level from charged current interactions while $B \to X_s\gamma$ and $B \to X_s\ l^+l^-$ first occur at one-loop. There are contributions to $B \to X_s\gamma$ and $B \to X_s\ l^+l^-$ from closed charm quark loops in addition to local operators in the effective theory resulting from integrating out the top quark and $W^\pm, Z^0$-bosons (we do not discuss up-quark contributions). It was recently suggested that these loop contributions give power corrections to the rate for $B \to X_s\gamma$ suppressed by the mass of the charm quark [6] as opposed to the mass of the bottom quark [4,5]. Explicit computation [6] showed that these new contributions are proportional to the chromomagnetic matrix element $\langle B|\overline{h}\sigma \cdot Gh|B\rangle$. Subsequently, it was shown [7,8] that such contributions to $B \to X_s\gamma$ do not dominate the $1/m_b^2$ power corrections found in [4,5]. In addition it was pointed out [7,8] that there is an infinite tower of operators whose contributions are suppressed only by powers of $m_b\Lambda_{\rm QCD}/m_c^2$ but which appear with numerically small coefficients.

The Feynman diagrams highlighted by Voloshin [6] yield a term that vanishes for on-shell photons in addition to the term modifying $B \to X_s\gamma$. In this work we determine the contribution of these diagrams to the inclusive decay $B \to X_s\ l^+l^-$ where $l$ denotes an electron or a muon. We do not re-examine the $\mathcal{O}\left(1/m_b^2\right)$ contributions [4,9] as we have nothing new to add. Further, we do not include the non-renormalization group pQCD corrections to the decay width [10]. Only the leading free-quark amplitude and the loop correction yielding this new non-perturbative contribution are considered.

For momentum scales much below the mass of the $W^{\pm}$ gauge boson it is appropriate to describe these rare decays by an effective hamiltonian

$$\mathcal{H}_{\rm eff} = \frac{4G_F}{\sqrt{2}} V_{bt}V_{st}^{\dagger} \sum_i \; c_i(\mu) \; \mathcal{O}_i(\mu) \quad , \tag{1}$$

where the $c_i(\mu)$ are Wilson coefficients of the operators $\mathcal{O}_i(\mu)$ renormalized at a scale $\mu$ [11–13]. We have assumed that $V_{bt}V_{st}^{\dagger} = -V_{bc}V_{sc}^{\dagger}$ in writing $\mathcal{H}_{\rm eff}$ this way and we assume that the leptons and strange quark are massless. In order to fix notation we list the operators (we use the same notation as [11])

$$\begin{aligned}
\mathcal{O}_1 &= \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\alpha} \; \overline{c}_{L,\beta}\gamma_{\mu}c_{L,\beta} \\
\mathcal{O}_2 &= \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\beta} \; \overline{c}_{L,\beta}\gamma_{\mu}c_{L,\alpha} \\
\mathcal{O}_3 &= \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\alpha} \; \left[\overline{u}_{L,\beta}\gamma_{\mu}u_{L,\beta} \; + \; ....\overline{b}_{L,\beta}\gamma_{\mu}b_{L,\beta}\right] \\
\mathcal{O}_4 &= \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\beta} \; \left[\overline{u}_{L,\beta}\gamma_{\mu}u_{L,\alpha} \; + \; ....\overline{b}_{L,\beta}\gamma_{\mu}b_{L,\alpha}\right] \\
\mathcal{O}_5 &= \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\alpha} \; \left[\overline{u}_{R,\beta}\gamma_{\mu}u_{R,\beta} \; + \; ....\overline{b}_{R,\beta}\gamma_{\mu}b_{R,\beta}\right] \\
\mathcal{O}_6 &= \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\beta} \; \left[\overline{u}_{R,\beta}\gamma_{\mu}u_{R,\alpha} \; + \; ....\overline{b}_{R,\beta}\gamma_{\mu}b_{R,\alpha}\right] \\
\mathcal{O}_7 &= \frac{e}{16\pi^2} m_b \overline{s}_{L,\alpha}\sigma^{\mu\nu}b_{R,\alpha} \; F_{\mu\nu} \\
\mathcal{O}_8 &= \frac{e^2}{16\pi^2} \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\alpha} \; \overline{l}\gamma_{\mu}l \\
\mathcal{O}_9 &= \frac{e^2}{16\pi^2} \overline{s}_{L,\alpha}\gamma^{\mu}b_{L,\alpha} \; \overline{l}\gamma_{\mu}\gamma_5 l \\
\mathcal{O}_{10} &= \frac{g}{16\pi^2} m_b \overline{s}_{L,\alpha}\sigma^{\mu\nu}T^a_{\alpha\beta}b_{R,\beta} \; G^a_{\mu\nu} \quad .
\end{aligned} \tag{2}$$

Working at leading order [11] we truncate the operator basis to $\mathcal{O}_1$, $\mathcal{O}_2$, $\mathcal{O}_7$, $\mathcal{O}_8$ and $\mathcal{O}_9$. The next to leading order results can be found in [12,13] and a complete summary including the non-perturbative and pQCD corrections can be found in [9].

The rate and other "short-distance" observables for the decay arise from an operator-product expansion of

$$\mathcal{T} = {\rm Im}\left( i \int \; d^4x \; e^{-iqx} \langle B| \; T\left[\mathcal{H}_{\rm eff}(x)\mathcal{H}^{\dagger}_{\rm eff}(0)\right] \; |B\rangle \right) \quad , \tag{3}$$

integrated over the appropriate leptonic phase space. The free quark decay result emerges as the leading term in the OPE of $\mathcal{T}$ with higher order corrections to the free quark result arising systematically [4,9] as a power series in $\alpha_s(\mu)$ and $1/m_b^n$. The component of $\mathcal{T}$ previously omitted from $B \to X_s\gamma$ [6] arises in

$$T\left[c_2\mathcal{O}_2(x) \; c_7^*\mathcal{O}_7^{\dagger}(0) \quad + \quad {\rm h.c.}\right] \quad , \tag{4}$$

where the charm loop closes off via a soft gluon and a photon, as shown in figure 1. The diagrams that modify $B \to X_s \; l^+l^-$ are shown in figure 2 (we have not shown the hermitian conjugate).

The amplitude resulting from closing the charm quark legs of $\mathcal{O}_2$ with the emission of a gluon of momentum $k$ with polarization $\eta$, and a photon of momentum $p$ with polarization $\lambda$ is given by (to leading order in $k$)

$$\mathcal{A} = \frac{e g_s Q_c}{4\pi^2}\ \overline{s} T^a \gamma^\mu (1-\gamma_5) b\ \left[ I_1\ \frac{1}{2} p_\eta \epsilon^{ab\lambda\mu} p_a k_b - I_1\ \frac{1}{2} p\cdot k \epsilon^{a\eta\lambda\mu} p_a + I_2\ p^2 \epsilon^{\lambda b\eta\mu} k_b - I_2\ p_\lambda \epsilon^{ab\eta\mu} p_a k_b \right] \quad , \quad (5)$$

where the functions $I_{1,2}$ are given by

$$I_{1,2} = \int_0^1\ dx \int_0^{1-x}\ dy \frac{2xy\ \ ,\ \ x(1-x)}{[m_c^2 - x(1-x)p^2 - 2xyp\cdot k]} \quad . \quad (6)$$

As $k^2 \sim \Lambda_{\rm QCD}^2 \ll m_c^2$ we neglected it in both the numerator of $\mathcal{A}$ and denominator of $I_{1,2}$. However the $p\cdot k$ term appearing in the denominator of $I_{1,2}$ is of order $\sim E_\gamma\ \Lambda_{\rm QCD} \sim m_c^2$ and dominates over the other terms near the charm pair threshold ($p^2 \sim 4m_c^2$) region. Before proceeding it is interesting to explore the issues raised in [7,8]. We expand the integrals $I_{1,2}$ in terms of $p\cdot k$ and evalute the coefficients of terms higher order in $\Lambda_{\rm QCD}$. After performing the angular integration over phase space one finds that

$$< I_1(s, m_c^2) > \ = \sum_{n=even}^{\infty}\ a_n\ b_n(s)\ \ (E_\gamma\ \Lambda_{\rm QCD})^n$$
$$b_n(s) = \int_0^1\ dx \frac{x^{n+1}(1-x)^{n+2}}{(m_c^2 - sx(1-x))^{n+1}} \quad , \quad (7)$$

where $E_\gamma$ is the energy of the lepton pair in the rest frame of the B-meson. The first few constants $a_n$ are $a_0 = \frac{1}{2}$ , $a_2 = \frac{2}{3}$ , $a_4 = \frac{16}{15}$. In the case of on-shell photons, $s = p^2 = 0$ and $E_\gamma = m_b/2$, one finds that the contribution from terms higher order in $\Lambda_{\rm QCD}$ converges rapidly. When the photon is off its mass-shell as is the case for $B \to X_s\ l^+ l^-$ and the invariant mass of the lepton pair are far from the charm threshold region the series is again convergent (requiring that $m_b - \sqrt{s} \gg \Lambda_{\rm QCD}$). In contrast, when the invariant mass of the lepton pair is near the charm threshold region the series is not convergent. This is to be expected since we know that the charmonium resonances in this region do not appear at any finite order in perturbation theory. In fact, this problem is not isolated to our computation but is a problem for the entire parton level computation of the differential rate. One expects that suitable smoothed quantities over this region of invariant mass will be reproduced by the same smoothed quantities in the parton model by duality. For the rest of our discussion we will drop the $p\cdot k$ terms so that the integrals $I_{1,2}$ become identical $I_1 = I_2 = I$ where

$$I(s, m_c^2) = \int_0^1\ dy\ \frac{y(1-y)^2}{[m_c^2 - y(1-y)s]} \quad . \quad (8)$$

The leading contribution to the differential decay rate comes from the free quark decay and at leading order in strong interactions [11] is

$$\frac{d\Gamma^{(0)}}{d\hat{s}} = \frac{\alpha^2}{4\pi^2}\frac{G_F^2 m_b^5}{192\pi^3}|V_{bt}V_{st}^\dagger|^2\ (1-\hat{s})^2\ \Big[ (1+2\hat{s})|c_8(m_b) + g(r,\hat{s})(c_2(m_b)+3c_1(m_b))|^2$$
$$+\ (1+2\hat{s})|c_9(m_b)|^2\ +\ 4(1+2/\hat{s})|c_7(m_b)|^2$$
$$+12{\rm Re}\left[c_7(m_b)^*(c_8(m_b) + g(r,\hat{s})(c_2(m_b)+3c_1(m_b)))\right]\Big] \quad , \quad (9)$$

where $r = m_c/m_b$ is the ratio of quark masses and $\hat{s} = p^2/m_b^2$ is the invariant mass of the lepton pair. The function $g(z,\hat{s})$ arises from closing the charm loop from operators $\mathcal{O}_{1,2}$ forming a virtual photon, and is given by (at $\mu = m_b$) [11–13]

$$g(z,\hat{s}) = -\frac{8}{9}lnz + \frac{8}{27} + \frac{4}{9}y \\ -\frac{2}{9}(2+y)\sqrt{|1-y|}\left[\theta(1-y)\left(ln\frac{1+\sqrt{1-y}}{1-\sqrt{1-y}} + i\pi\right) + \theta(y-1)2\tan^{-1}\frac{1}{\sqrt{y-1}}\right] \quad . \quad (10)$$

with $y = \frac{4z^2}{\hat{s}}$.

Computation of the non-perturbative term is straightforward but tedious and we will not detail it here. The additional contribution to the differential rate is found to be

$$\frac{d\Gamma}{d\hat{s}} = \frac{\alpha^2}{4\pi^2}\,\frac{G_F^2 m_b^5}{192\pi^3}\,|V_{bt}V_{st}^\dagger|^2\,\frac{32\lambda_2}{3m_b^2}\,(1-\hat{s})^2 \\ \mathrm{Re}\,[(\ (1/\hat{s}+6-\hat{s})\ c_7(m_b)\ +\ (2+\hat{s})\ (c_8(m_b)\ +\ (c_2(m_b)\ +\ 3\ c_1(m_b))\ g(r,\hat{s})\ )\ )^* \\ c_2(m_b)\ I(\hat{s},r)\ ] \quad . \quad (11)$$

The nonperturbative matrix element $\lambda_2$ that appears in this contribution is found from the $B^* - B$ mass difference to be $\lambda_2 \sim 0.1\mathrm{GeV}^2$. Comparing the size of this new non-perturbative contribution to the differential decay rate, as shown in figure 3, one see that it is small except near the charm threshold. Integrating the expression for the rates between $s = m_b^2/2$ and $s = m_b^2$ we find the rate to be decreased by $\sim 1.5\%$ due to this new contribution (i.e. smaller than the contribution from the $1/m_b^2$ operators, $\sim 3\%$). Unfortunately near threshold is precisely the region where the above computation fails due to the presense of incalculable long-distance physics associated with the charmonium resonances. In the region either far above or far below the charm threshold the correction considered here can be safely neglected.

In conclusion, we have determined a previously omitted non-perturbative contribution to $B \to X_s\ l^+l^-$, that is not explicitly suppressed by $1/m_b^n$. This has the same origin as the $1/m_c^2$ contribution to $B \to X_s\gamma$ recently observed by Voloshin [6]. We find that in the regions of lepton invariant mass where we can reliably compute such contributions they are small. The computed amplitude has interesting threshold behavior near $s = 4m_c^2$ but unfortunately large strong interactions corrections that we cannot determine will significantly modify the result in this region.

## ACKNOWLEDGMENTS

This work is partially supported by the US Department of Energy.




FIG. 1. A contribution to $B \to X_s\gamma$ from the time-ordered product $T[\ \mathcal{H}_{\rm eff}(x)\ \mathcal{H}^\dagger_{\rm eff}(0)\ ]$ proportional to $\lambda_2$ and naively suppressed by $1/m_c^2$.

## REFERENCES


[1] R. Balest *et al.*, (CLEO Collaboration), presented at "International Conference on High Energy Physics", Glasgow, Scotland (1994), edited by P.J. Bussey and I.G. Knowles.

[2] M. Voloshin and M. Shifman, Yad. Fiz. 41 (1985) 187; J. Chay, H. Georgi and B. Grinstein, Phys. Lett. **B247** (1990) 399.

[3] I.I. Bigi, N.G. Uraltsev and A.I. Vainshtein, Phys. Lett. **B293** (1992) 430; Phys. Lett. **B297** (1993) 477 (E); I.I. Bigi et al., Phys. Rev. Lett. **71** (1993) 496; A.V. Manohar and M.B. Wise, Phys. Rev. **D49** (1994) 1310.

[4] A. Falk, M. Luke and M.J. Savage, Phys. Rev. **D49** (1994) 3367.

[5] I.I. Bigi, N.G. Uraltsev and A.I. Vainshtein, Phys. Lett. **B247** (1992) 430.

[6] M. B. Voloshin, Phys. Lett. **B397** (1997) 275.

[7] Z. Ligeti, L. Randall and M.B. Wise, Caltech Preprint CALT-68-2097 (1997), hep-ph/9702322.

[8] A.K. Grant, A.G. Morgan, S. Nussinov and R.D. Peccei, UCLA Preprint UCLA/97/TEP/5 (1997).

[9] A. Ali, G. Hiller, L.T. Handoko and T. Morozumi, Phys. Rev. **D55** (1997) 4105.

[10] A. Ali and c. Greub, Z. Phys. **C49** (1991) 431; Phys. Lett. **B259** (1991) 182; Z. Phys. **C60** (1993) 433; Phys. Lett. **B361** (1995) 146; N. Pott, Phys. Rev **D54** (1996) 938.

[11] B. Grinstein, M.J. Savage and M.B. Wise, Nucl. Phys. **B319** (1989) 271.

[12] M. Misiak, Nucl. Phys. **B393** (1993) 23; Nucl. Phys. **B439** (1995) 461 (E);

[13] A.J. Buras and M. Munz, Phys. Rev. **D52** (1995) 186.




FIG. 2. Contributions to $B \to X_s\ l^+l^-$ from the time-ordered product $T[\ \mathcal{H}_{\rm eff}(x)\ \mathcal{H}^\dagger_{\rm eff}(0)\ ]$ proportional to $\lambda_2$.




FIG. 3. The differential decay rate for inclusive $B \to X_s\, l^+ l^-$ using quark masses $m_c = 1.5$ GeV and $m_b = 4.5$ GeV normalized to the semileptonic width. The dashed line is the leading order, free-quark contribution and the solid line is the sum of the free-quark and the non-perturbative contribution computed in this work. Inset is the distribution in the charm threshold region. We have used $c_1 = 0.22$, $c_2 = -1.09$, $c_7 = 0.31$, $c_8 = -4.16$ and $c_9 = 4.87$ at the scale $\mu = m_b$.